

# *Past, Present and Future of Near-Infrared Imaging*

**BRITTON CHANCE, SHOKO NIOKA, JUN ZHANG,  
XAVIER INTES, YU CHEN, CHENPENG MU,  
DANA BLESSINGTON, LANLAN ZHOU,  
AND ZHIHONG ZHANG**

---

**DEPARTMENT OF BIOCHEMISTRY AND BIOPHYSICS and RADIOLOGY  
UNIVERSITY OF PENNSYLVANIA**

**Support: NIH Grants CA878046, CA27895, and CO97065**



## Sensitivity/Specificity

a PPV, NPV >90%

## Tumor/Tissue Ratio

## Intrinsic Signals

### Angiogenesis

Blood volume  
ICG Occupancy

~ 2  
~ 3

### Hypermetabolism

Deoxygenation of hemoglobin  
Increased Fluorescence of Glucose

2  
> 3

### Extrinsic Probes


Blood Pool -- ICG, FDA approved

2-3

### NIR - Molecular Beacons

Targeted cytochrome, LDL, Cathepsin B  
(ordinary, stealth)

>10



BC 368b

Chance '03




# *Amplitude Cancellation System*

## *Human Breast Test*


(AS:24 12/10/97)

**Right breast – model**




**Blood  
Deoxygenation**

**Blood  
Volume**




**Left breast – model**



**Blood  
Deoxygenation**

**Blood  
Volume**




SNH 60

Chance '03



# Dual Wavelength Amplitude Cancellation (AC-TM) Imager


## 2D Angiogenesis vs. Hypermetabolism Imagery, 71 subjects (15 cancers)




	Zone 1	Zone II		Zone 1	Zone II
Sensitivity	91.7%	100%	PPV	100%	70.9%
Specificity	100%	90.6%	NPV	98.1%	100%

# Analysis of 71 Subjects with Continuous Wave Near Infrared Image


Receiver Operator Curve of Blood Volume




Receiver Operator Curve of Deoxygenation




Receiver Operator Curve of Blood Volume and Deoxygenation




Breast Test Imaging



# *The possibilities of nanotechnology*



# NADH signal of Rif-1 Tumor



**\*Depths: 200um to 3100um below skin surface**





# 3D Fluorescent Images of c-MYC Induced Mammary Tumor



Surface Imaging  
(Outline)

Volume Rendering Imaging  
(Fluorescent Intensity)

PN



Pyro-2DG



3D fluorescence images of c-MYC induced mammary tumor reflect the outline and the metabolic state of the tumor. These 3D images were reconstructed by 9 slices of fluorescent signal from the top to the bottom of the tumor. The size of each frame was  $1.024 \text{ cm} \times 1.024 \text{ cm}$ .






# HepG2 tumor + 0.2ul 1mg/ml Pyro-2DG after 2hrs (#M11)

Fp




PN

Pyro-2DG

Top




Bottom




# 9L glioma + Pyro-2DG after 2hrs (Top Volume View)





**Fp**



**PN**



**Pyro-2DG**





# Quantitative evaluation of Cy5.5 labeled Cathepsin B beacon delivery by 3D high resolution imaging



Direct Volume Rendering

Surface Rendering

Top  
view



Bottom  
view



3D image of stealth beacon detection of animal cancer.  
Note island of dissemination of cancer.

1 cm x 1 cm  
Depth: 200 ~ 1000  $\mu$ m